

ОРИГИНАЛНИ НАУЧНИ ЧЛАНЦИ
ОРИГИНАЛЬНЫЕ НАУЧНЫЕ СТАТЬИ
ORIGINAL SCIENTIFIC PAPERS

SOME CRITICAL REMARKS
ON THE PAPER "A NOTE ON THE
METRIZABILITY OF TVS-CONE
METRIC SPACES"

Suzana M. Aleksić^a, Ljiljana R. Paunović^b, Stojan N. Radenović^c,
Francesca Vetro^d

^a University of Kragujevac, Faculty of Science, Department of Mathematics and Informatics, Kragujevac, Republic of Serbia,
e-mail: suzanasimic@kg.ac.rs,
ORCID iD: <http://orcid.org/0000-0002-2176-5091>,

^b University of Priština - Kosovska Mitrovica, Faculty of Education in Prizren - Leposavić, Leposavić, Republic of Serbia,
e-mail: ljiljana.paunovic76@gmail.com,
ORCID iD: <http://orcid.org/0000-0002-5449-9367>.

^c University of Belgrade, Faculty of Mechanical Engineering, Belgrade, Republic of Serbia,
e-mail: radens@beotel.net,
ORCID iD: <http://orcid.org/0000-0001-8254-6688>

^d University of Palermo, Department of Energy, Information Engineering and Mathematical Models (DEIM), Palermo, Italy,
e-mail: francesca.vetro@unipa.it,
ORCID iD: <http://orcid.org/0000-0001-7448-5299>

<http://dx.doi.org/10.5937/vojtehg66-15128>

FIELD: Mathematics

ARTICLE TYPE: Original Scientific Paper

ARTICLE LANGUAGE: English

Abstract:

This short and concise note provides a detailed exposition of the approach and results established by (Lin et al, 2015, pp.271-279). We show that the obtained results are not particularly surprising and new. Namely, using an old result due to K. Deimling it is indicated that tvs-cone metric spaces over

ACKNOWLEDGMENT: The first author is grateful for the financial support from the Ministry of Education, Science and Technological Development of the Republic of Serbia (Grant number #174024). The second author is grateful for the financial support from the Ministry of Education, Science and Technological Development of the Republic of Serbia (Grant number #174002).

solid cones are actually cone metric spaces over normal solid cones. Hence, there are only cone metric spaces over normal solid cones or over normal non-solid cones. One question still unanswered is whether an ordered topological vector space with a non-normal non-solid cone exists.

Key words: tvs-cone metric space, metrizable, solid, normal, non-normal.

Introduction, preliminaries and results

We will denote by E a topological vector space with the zero vector θ . A subset C of E is called a cone in E if and only if:

- (a) C is nonempty and closed in E ,
- (b) $x, y \in C$ and $a, b \in R^+$ imply $ax + by \in C$,
- (c) $x, -x \in C$ implies $x = \theta$.

For more details we refer the reader to (Ansari et al, 2016), (Deimling, 1985), (Filipović et al, 2011), (Janković et al, 2011), (Köthe, 1969), (Shaefer, 1971) and (Wong & Ng, 1973).

For a given cone C , a partial ordering \leq with respect to C is introduced in the following way: $x \leq y$ if and only if $y - x \in C$. We write $x < y$ to indicate that $x \leq y$, but $x \neq y$. If $y - x \in \text{int } C$, we write $x \ll y$. It is clear that $<$ and \ll are not relations of partial order.

The pair (E, C) is called an ordered topological vector space. Throughout the paper, C stands for a solid cone, i.e. $\text{int } C \neq \emptyset$. In most papers it turns out that this assumption is essential (Amini-Harandi & Fakhar, 2010), (Cakalli et al, 2012), (Du, 2010), (Proinov, 2013), (Khani & Pourmahdian, 2011), (Lin et al, 2015), (Simić, 2011), (Vandergraft, 1967) and (Zabrejko, 1997).

The following result from (Wong & Ng, 1973) is important in the framework of ordered topological vector spaces.

Proposition 1.1. Let (E, C) be an ordered topological vector space.

Then e is an interior point of C , i.e., $e \in \text{int } C$, if and only if

$$[-e, e] = \{x \in E : -e \leq x \leq e\} = (e - C) \cap (C - e)$$

is the neighborhood of θ .

Under the assumption $E \neq \{\theta\}$, we have the following corollary.

Corollary 1.2. Let (E, C) be an ordered topological vector space with $\text{int } C \neq \emptyset$. Then

- (a) $\theta \notin \text{int } C$,
- (b) $\lambda \text{int } C = \text{int } C$ if $\lambda > 0$ and $C + \text{int } C = \text{int } C$,
- (c) (E, C) is Hausdorff ordered topological vector space.

Let us note that the property (c) is an immediate consequence of Theorem 2.1 (2) from (Jachymski & Klima, 2016).

For each $c \in \text{int } C$ there exists a norm $\|\cdot\|_c$ on the vector space E where $\|\cdot\|_c$ is the Minkowski functional on $E = E_c = \bigcup_{n \in \mathbb{N}} n[-c, c]$, i.e.,

$$\|x\|_c = \inf \{\lambda > 0 : x \in \lambda[-c, c]\}.$$

It is obvious that all norms $\|\cdot\|_c$, $c \in \text{int } C$, are equivalent. Further, we get that $(E, \|\cdot\|_c, C)$ is an ordered normed space with the normal-solid cone C . For more details see (Deimling, 1985, pp.230, Proposition 19.9). Therefore, $\text{int } C$ under the topology on E is a subset of $\text{int } C$ under the norm topology, i.e., $(\text{int } C)_E \subseteq (\text{int } C)_{\|\cdot\|_c}$. However, the converse is also valid. Hence, $(\text{int } C)_E = (\text{int } C)_{\|\cdot\|_c}$.

From what has already been proved, it is clear that the notion of tvs-cone metric space in the sense of Du, see (Du, 2010), is the same as that of (Lin et al, 2015, Definition 2.3).

According to (Deimling, 1985) and (Kadelburg et al, 2016), we are thus led to the following strengthening conclusions.

Proposition 1.3. Each tvs-cone metric space over a solid cone in the sense of the both definitions (Du, 2010) and (Lin et al, 2015) is in fact a cone metric space over a normal solid cone. Therefore, the well known results, (Amini-Harandi & Fakhar, 2010), (Du, 2010), (Kadelburg et al, 2011), (Khani & Pourmahdian, 2011) give a positive answer to Question 1.1 that arises in (Lin et al, 2015).

Proposition 1.4. The claims from (Lin, 2015) are the immediate consequences of the corresponding results in the framework of cone metric spaces over normal solid cones, see (Huang & Zhang, 2007), (Janković et al, 2011), (Kadelburg et al, 2011), (Simić, 2011).

Proposition 1.5. If C is a solid cone in some ordered topological vector space E , then it is a normal solid cone under a suitable defined norm. For all details see (Deimling, 1985, Proposition 19.9).

Remark 1.6. It is worth pointing out that in (Du, 2010), (Fierro, 2016), (Kadelburg et al, 2016), (Lin et al, 2015) and (Simić, 2011) the assumption:

(E, C) is Hausdorff, is superfluous. Applying Corollary 1.2 (c) we conclude that the condition $\text{int } C \neq \emptyset$ implies that the ordered topological vector space (E, C) is Hausdorff.

In the theory of abstract metric spaces, quite a few proofs are based on c-sequences introduced in (Alnafesi et al, 2011). Namely, for a sequence $\{x_n\}$ in the solid cone C of a real ordered topological vector space E , we say that it is a c-sequence if for every interior point c of the cone C there exists $n_0 \in \mathbb{N}$ such that $x_n << c$ whenever $n \geq n_0$. If C is normal in E , then $\{x_n\}$ is a c-sequence if and only if $x_n \rightarrow \theta$ in E . However, if the cone C is a non-normal solid cone, then the previous is not true, i.e., a c-sequence does not always converge to θ as $n \rightarrow \infty$. We refer the readers to (Radenović et al, 2017), (Xu & Radenović, 2014), (Xu et al, 2016), (Đorđević et al, 2011) and (Kadelburg et al, 2011) for more details on c-sequences.

Theorem 1.7. Let $(E, \|\cdot\|)$ be an ordered (real) Banach space with the underlying solid cone C and $\{x_n\}$ be a sequence in C . Then $\{x_n\}$ is a c-sequence if and only if $\|x_n\|_e \rightarrow \theta$ as $n \rightarrow \infty$ where

$$\|x\|_e = \inf\{\lambda > 0 : x \in \lambda[-e, e], e \in \text{int } C\},$$

i.e., $\|\cdot\|_e$ is the Minkowski functional.

Proof. First, $\|\cdot\|_e$ is a norm in E . If C is a normal solid cone, then this

norm is equivalent to the given one, and $\|x_n\| \rightarrow \theta$ if and only if $\|x_n\|_e \rightarrow \theta$.

If C is not normal, then C is a normal and solid cone under the norm $\|\cdot\|_e$ for each $e \in \text{int } C$. Since $\|\cdot\|$ and $\|\cdot\|_e$ give the same interior points, then by the first case $\{x_n\}$ is a c-sequence if and only if $\|x_n\|_e \rightarrow \theta$. Consequently, in each normal cone, the conditions $x_n \rightarrow \theta$ and $\{x_n\}$ is a c-sequence are equivalent.

Conclusion

Note that we have actually proved that there exist cone metric spaces over normal solid cones or over normal non-solid cones. Therefore, the

results obtained in (Lin et al, 2015) are not new but rather the consequences of known facts (Huang & Zhang, 2007), (Janković et al, 2011), (Kadelburg et al, 2011) and (Simić, 2011). We can conclude that the existence of ordered topological vector spaces with a non-normal non-solid cone is still an open question.

References

- Alnafesi, S.H., Radenović, S., & Shahzad, N., 2011. Fixed point theorems for mappings with convex diminishing diameters on cone metric spaces. *Appl. Math. Lett.*, 24 (2), pp.2162-2166. Available at:<https://doi.org/10.1016/j.aml.2011.06.019>.
- Amini-Harandi, A., & Fakhar, M., 2010. Fixed point theory in cone metric spaces obtained via the scalarization method. *Comput. Math. Appl.*, 59 (11), pp.3529-3534. Available at: <https://doi.org/10.1016/j.camwa.2010.03.046>.
- Ansari, A.H., Chandok, S., Hussain, N., & Paunović, L., 2016. Fixed point of weak contractions in regular cone. *Journal of Advanced Mathematical Studies*, 9 (1), pp.72-82.
- Deimling, K., 1985. *Nonlinear Functional Analysis*. Springer-Verlag.
- Du, W.S., 2010. A note on cone metric fixed point theory and its equivalence. *Nonlinear Anal.*, 72 (5), pp.2259-2261. Available at: <https://doi.org/10.1016/j.na.2009.10.026>.
- Đorđević, M., Đorić, D., Kadelburg, Z., Radenović, S., & Spasić, D., 2011. Fixed point results under C-distance in TVS- cone metric spaces. *Fixed Point Theory and Applications*, 2011:29. Available at: <https://doi.org/10.1186/1687-1812-2011-29>.
- Filipović, M., Paunović, Lj., Radenović, S., & Rajović, M., 2011. Remarks on "Cone Metric Spaces and Fixed Point Theorems of T-Kannan Contractive Mappings". *Mathematical and Computer Modelling*, 54 (5-6). Available at: <https://doi.org/10.1016/j.mcm.2011.04.018>.
- Huang, L.G., & Zhang, X., 2007. Cone metric spaces and fixed point theorems of contractive mappings. *J. Math. Anal. Appl.*, 332 (2), pp.1468-1476. Available at: <https://doi.org/10.1016/j.jmaa.2005.03.087>.
- Jachymski, J., & Klima, J., 2016. Cantor's intersection theorem for K-metric spaces with a solid cone and a contraction principle. *J. Fixed Point Theory Appl.*, 18 (3), pp.445-463. Available at: <https://doi.org/10.1007/s11784-016-0312-1>.
- Janković, S., Kadelburg, Z., & Radenović, S., 2011. On cone metric spaces: A survey. *Nonlinear Anal.*, 74 (7), pp.2591-2601. Available at: <https://doi.org/10.1016/j.na.2010.12.014>.

- Kadelburg, Z., Paunović, L., Radenović, S., & Rad, G.S., 2016. Non-normal cone metric and cone b-metric spaces and fixed point results. *Scientific publications of the state University of Novi Pazar, Ser. A: Appl. Math. Inform. And Mech.*, 8 (2), pp.177-186.
Available at: <https://doi.org/10.5937/SPSUNP1602177K>.
- Kadelburg, Z., Radenović, S., & Rakočević, V., 2011. A note on the equivalence of some metric and cone metric fixed point results. *Appl. Math. Lett.*, 24 (3), pp.370-374. Available at: <https://doi.org/10.1016/j.aml.2010.10.030>.
- Khani, M., & Pourmahdian, M., 2011. On the metrizability of cone metric space. *Topology Appl.*, 158 (2), pp.190-193. Available at: <https://doi.org/10.1016/j.topol.2010.10.016>.
- Köthe, G., 1969. *Topological Vector Spaces I*. New York: Springer-Verlag, Inc..
- Lin, S., Li, K., & Ge, Y., 2015. On the metrizability of tvs-cone metric spaces, Publication de l'Institut Mathématique. Nouvelle série, tome, 98 (112), pp.271-279. Available at: <https://doi.org/10.2298/PIM1512271L>.
- Radenović, S., Vetro, F., & Xu, S., 2017. Some new results on perov type mappings. *J. Adv. Math. Stud.*, 10(3), pp.396-409.
- Shaefer, H.H., 1971. *Topological Vector Spaces*, 3rd ed. New York: Springer.
- Simić, S., 2011. A note on Stone's, Baire's, Ky Fan's and Dugundji's theorem in tvs-cone metric spaces. *Appl. Math. Lett.*, 24 (6), pp.999-1002. Available at: <https://doi.org/10.1016/j.aml.2011.01.014>.
- Vandergraft, J.S., 1967. Newton's method for convex operators in partially ordered spaces. *SIAM J. Num. Anal.*, 4 (3), pp.406-432. Available at: <https://doi.org/10.1137/0704037>.
- Wong, Y.C., & Ng, K.F., 1973. *Partially Ordered Topological Vector Spaces*. Oxford: Clarendon Press.
- Xu, S., & Radenović, S., 2014. Fixed point theorems of generalized Lipschitz mappings on cone metric spaces over Banach algebras without assumption of normality. *Fixed Point Theory Appl.*, 2014:102. Available at: <https://doi.org/10.1186/1687-1812-2014-102>.
- Xu, S., Doličanin, Č., & Radenović, S., 2016. Some remarks on results of Perov type. *J. Adv. Math. Stud.*, 9 (3), pp.361-369.
- Zabrejko, P.P., 1997. K-metric and K-normed linear spaces, survey. *Collect. Math.*, 48, pp.825-859.

НЕКОТОРЫЕ КРИТИЧЕСКИЕ ЗАМЕЧАНИЯ О РАБОТЕ «ЗАМЕТКИ О МЕТРИЗУЕМОСТИ ТВП-КОНИЧЕСКИХ ПРОСТРАНСТВ»

Сузана М. Алексич^a, Лилияна Р. Паунович^b, Стоян Н. Раденович^b,
Франческа Ветро^г

^a Университет в г. Крагуевац, Естественно-математический факультет,
Институт математики и информатики, г. Крагуевац, Республика Сербия,

^b Университет в Приштине–Косовской Митровице, Педагогический
факультет в Призрене–Лепосавиче, г. Лепосавич, Республика Сербия,

^в Белградский университет, Машиностроительный факультет, г. Белград,
Республика Сербия,

^г Университет в г. Палермо, Департамент энергетики, информационного
инжиниринга и математического моделирования, г. Палермо,
Итальянская Республика

ОБЛАСТЬ: математика

ВИД СТАТЬИ: оригинальная научная статья

ЯЗЫК СТАТЬИ, английский

Резюме:

В представленных, в данной статье, заметках приведен подробный обзор методов и полученных результатов исследования группы авторов, во главе с Шой Линь (Lin et al, 2015, pp.271-279). Мы в свою очередь доказываем, что их результаты не являются инновационными. В частности, при применении известных результатов К. Деймлинга установлено, что ТВП-конические метрические пространства с конусами с непустой внутренностью фактически являются метрическими пространствами с нормальной конической и непустой внутренностью. Следовательно, существуют только конические метрические пространства с нормальными конусами, внутренность которых непуста или конусами, которые нормальны, но с пустой внутренностью. До сих пор не установлено упорядоченное топологическое векторное пространство с конусом, который не является нормальным, а внутренность которого не пуста.

Ключевые слова: твп-коническое метрическое пространство, метризуемость, конус с непустой внутренностью, нормальный, ненормальный.

НЕКЕ КРИТИЧКЕ НАПОМЕНЕ О РАДУ „БЕЛЕШКА О МЕТРИЗАБИЛНОСТИ ТВП-КОНУСНИХ МЕТРИЧКИХ ПРОСТОРА”

Сузана М. Алексић^a, Љиљана Р. Паунович^b, Стојан Н. Раденовић^b,
Франческа Ветро^г

^a Универзитет у Крагујевцу, Природно-математички факултет,
Институт за математику и информатику, Крагујевац, Република Србија,

- ⁶ Универзитет у Приштини – Косовској Митровици, Учитељски факултет у Призрену – Лепосавићу, Лепосавић, Република Србија
^в Универзитет у Београду, Машински факултет, Београд, Република Србија
^г Универзитет у Палерму, Одсек за енергетику, информационо инжењерство и математичке моделе, Палермо, Италија

ОБЛАСТ: математика
ВРСТА ЧЛАНКА: оригинални научни чланак
ЈЕЗИК ЧЛАНКА: енглески

Сажетак:

Ова кратка и преугледна белешка даје детаљан извештај о приступу и резултатима до којих су дошли Шоу Лин и група аутора (*Lin et al., 2015, pp.271-279*). У чланку је показано да њихови резултати нису нарочито изненађујући и нови. У ствари, коришћењем једног познатог К. Демлинговог резултата назначено је да су твр-конусни метрички простори са конусима који имају непразну унутрашњост заправо конусни метрички простори са нормалним конусима и непразним унутрашњостима. Стoga, постоје само конусни метрички простори са нормалним конусима чија унутрашњост није празна или са конусима који су нормални, али са празним унутрашњостима. Још увек се не зна да ли постоји уређен тополошки векторски простор са конусом који није нормалан и чија унутрашњост није празна.

Кључне речи: твр-конусни метрички простор, метризабилан, конус са непразном унутрашњошћу, нормалан, није нормалан.

Paper received on / Дата получения работы / Датум пријема чланка: 26.09.2017.
Manuscript corrections submitted on / Дата получения исправленной версии работы /
Датум достављања исправки рукописа: 10.10.2017.
Paper accepted for publishing on / Дата окончательного согласования работы / Датум
коначног прихватања члана за објављивање: 12.10.2017.

© 2018 The Authors. Published by Vojnotehnički glasnik / Military Technical Courier (www.vtg.mod.gov.rs, втг.мо.упр.срб). This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/rs/>).

© 2018 Авторы. Опубликовано в «Военно-технический вестник / Vojnotehnički glasnik / Military Technical Courier» (www.vtg.mod.gov.rs, втг.мо.упр.срб). Данная статья в открытом доступе и распространяется в соответствии с лицензией «Creative Commons» (<http://creativecommons.org/licenses/by/3.0/rs/>).

© 2018 Аутори. Објавио Војнотехнички гласник / Vojnotehnički glasnik / Military Technical Courier (www.vtg.mod.gov.rs, втг.мо.упр.срб). Ово је чланак отвореног приступа и дистрибуира се у складу са Creative Commons licencem (<http://creativecommons.org/licenses/by/3.0/rs/>).

